

Contents

| | |
|---|---|
| <u>Definition</u> | 2 |
| <u>Prevalence and incidence</u> | 2 |
| <u>Signs and symptoms</u> | 2 |
| <u>Causes/risk factors</u> | 3 |
| <u>Pathophysiology (mechanism of disease)</u> | 4 |
| <u>Prognosis and complications</u> | 4 |
| <u>Diagnosis/detection</u> | 5 |
| <u>Parity of esteem</u> | 5 |
| <u>Pharmacological treatment</u> | 6 |
| <u>Non-pharmacological treatment</u> | 7 |
| <u>Patient support</u> | 8 |
| <u>Further resources</u> | 8 |
| <u>External websites</u> | 9 |
| <u>References</u> | 9 |

Definition

Bipolar disorder (also known as bipolar affective disorder) is a serious, long-term mental illness usually characterised by episodes of depressed mood and episodes of elated mood and increased activity; these are referred to as mania or hypomania (a milder form of mania).¹ Bipolar disorder has previously been known as manic depression, a term which is no longer used.²

The *Diagnostic and statistical manual of mental disorders 5th edition* (DSM-5) distinguishes between bipolar I disorder and bipolar II disorder.

- Bipolar I disorder is characterised by at least one manic episode with or without a history of major depressive episodes.
- Bipolar II disorder is characterised by one or more major depressive episodes and by at least one hypomanic episode, but no evidence of mania.¹

For information on how manic, hypomanic and depressive episodes are defined, and rapid-cycling bipolar disorder, visit the NICE Clinical Knowledge Summary (CKS) page ***Bipolar disorder – Definition, What is it?***

[Return to contents](#)

Prevalence and incidence

The **NICE guidance [CG185]** states that *‘the most widely accepted estimates of lifetime prevalence in adults of bipolar disorder (mania and depression) is 1% and bipolar II disorder (hypomania and depression) is 0.4%’*

Bipolar disorder can occur at any age, although it often develops between the ages of 15 and 19. Men and women from all backgrounds are equally likely to develop bipolar disorder.²

[Return to contents](#)

Signs and symptoms

The NHS website summarises the symptoms of the depression and mania associated with bipolar disorder in the following way:

| Symptoms of depression | Symptoms of mania |
|---|--|
| <ul style="list-style-type: none"> • <i>‘feeling sad, hopeless or irritable most of the time</i> • <i>lacking energy</i> • <i>difficulty concentrating and remembering things</i> • <i>loss of interest in everyday activities</i> • <i>feelings of emptiness or worthlessness</i> • <i>feelings of guilt and despair</i> • <i>feeling pessimistic about everything</i> • <i>self-doubt</i> | <ul style="list-style-type: none"> • <i>‘feeling very happy, elated or overjoyed</i> • <i>talking very quickly</i> • <i>feeling full of energy</i> • <i>feeling self-important/grandiosity</i> • <i>feeling full of great new ideas and having important plans</i> • <i>being easily distracted</i> • <i>being easily irritated or agitated</i> • <i>being delusional, having hallucinations and disturbed or illogical thinking</i> |

- | | |
|---|--|
| <ul style="list-style-type: none">• <i>being delusional, having hallucinations and disturbed or illogical thinking</i>• <i>lack of appetite</i>• <i>difficulty sleeping</i>• <i>waking up early</i>• <i>suicidal thoughts</i>³ | <ul style="list-style-type: none">• <i>not feeling like sleeping</i>• <i>not eating</i>• <i>doing things that often have disastrous consequences – such as spending large sums of money on expensive and sometimes unaffordable items</i>• <i>making decisions or saying things that are out of character and that others see as being risky or harmful</i>³ |
|---|--|

Watch Mind's video ***Bipolar Disorder, Talking about mental health – Episode 7*** to hear Laura, Steve and Joe talking about living with bipolar disorder.



[Return to contents](#)

Causes/risk factors

There is likely a combination of factors that can increase the risk of developing bipolar disorder or precipitate it. Suggested factors which increase risk include:

Psychosocial influences

- Childhood trauma, eg, physical or sexual abuse, neglect or bereavement.
- Stressful life events, eg, relationship breakdown, money worries and poverty, or experiencing a traumatic loss.
- Social exclusion.^{4,5}

Brain chemistry

- Dysfunction of the hypothalamic-pituitary-adrenal axis (HPA axis – a complex network of signals between the hypothalamus, the pituitary gland and the adrenal cortex) with abnormal secretion of cortisol.

- Neurotransmitters are thought to play a role due to medicines acting on them being shown to treat bipolar symptoms.^{4,5}

Genetic inheritance

It is thought that there is a significant genetic component to bipolar disorder. Those with a close family member with bipolar disorder have a five to ten-time greater lifetime risk of developing the illness when compared to the general population.^{4,5}

It should be noted that although prescribed medicines, illicit drugs and alcohol do not cause bipolar disorder, they can cause bipolar moods and symptoms.⁵ More information can be found on the Mind **What causes bipolar disorder?** page.

[Return to contents](#)

Pathophysiology (mechanism of disease)

The pathophysiology of bipolar disorder is still not well understood, despite extensive research. The following article explores the potential causes of bipolar disorder based on brain neuroimaging, genetics, neurophysiological, neurochemical and neuroimmuno-endocrinological approaches.

Sigitova E, et al. **Biological hypotheses and biomarkers of bipolar disorder**. *Psychiatry and Clinical Neurosciences* 2017; 71: 77–103.

[Return to contents](#)

Prognosis and complications

The NICE Clinical Knowledge Summary (CKS) *Bipolar disorder* states that, 'For most people, bipolar disorder is a chronic, lifelong illness, with an estimated average of 10 episodes experienced in a person's lifetime.

The risk of recurrence in the 12 months after an episode is especially high compared with other psychiatric disorders (50 percent at one year and 75 percent at four years).

*Recovery may or may not be complete between episodes. Incomplete recovery leads to an increased risk of relapse, greater functional impairment, and reduced quality of life.*⁶

Suicide and deliberate self-harm are the most significant complications of bipolar disorder. Severe, untreated bipolar disorder is associated with a 15 percent lifetime risk for suicide.⁷ It is estimated that 25 to 56 percent of people with bipolar disorder present with at least one suicide attempt in their lifetime and 14 to 59 percent have suicidal ideation.⁸ Suicide risk is higher earlier in the course of the illness at around 25 percent.⁷

Acute episodes of bipolar disorder can result in difficulty with finances and potentially financial ruin, traumatic injury, sexually transmitted infections and unplanned pregnancy, damage to reputation, occupation, and relationships, self-neglect, exhaustion, and dehydration, exploitation, alcohol and

substance misuse and reduced quality of life.⁸ There may also be harm to others as a result of neglect, depressive or paranoid delusions, grandiosity (an exaggerated belief of importance), overspending, poor judgement, and erratic or chaotic behaviour and, rarely, violence and aggression.⁸

People with bipolar disorder have an increased risk of cardiovascular disease and are twice as likely to die from it compared to the general population.⁸ There are a range of factors that are thought to contribute to this including poor diet and smoking, adverse effects of antipsychotic drugs, such as weight gain, alcohol and illegal drug misuse, reduced use of cardiovascular drugs (for example statins), and reduced likelihood to receive hospital care or operations such as stents. Additionally, those with bipolar disease are at increased risk of chronic kidney disease (CKD), respiratory disease and diabetes.⁸

[Return to contents](#)

Diagnosis/detection

Bipolar disorder may be suspected in those who display a mixture of the symptoms listed above. At this point they should be referred to a specialist for assessment.

NICE clinical guideline ***Bipolar disorder: assessment and management [CG185]*** states that, '*when adults present in primary care with depression, ask about previous periods of overactivity or disinhibited behaviour. If the overactivity or disinhibited behaviour lasted for four days or more, consider referral for a specialist mental health assessment... Refer people urgently for a specialist mental health assessment if mania or severe depression is suspected or they are a danger to themselves or others... Do not use questionnaires in primary care to identify bipolar disorder in adults.*'⁹

And in relation to children and young people, '*diagnosis of bipolar disorder in children or young people should be made only after a period of intensive, prospective longitudinal monitoring by a healthcare professional or multidisciplinary team trained and experienced in the assessment, diagnosis and management of bipolar disorder in children and young people, and in collaboration with the child or young person's parents or carers.*'⁹

NICE recommendations for assessing for bipolar disorder in secondary care include undertaking a full psychiatric assessment and establishing a person's history in terms of their mood, manic and depressive episodes, social history, comorbidities, family history and previous treatment.⁹

For more information on diagnosis access NICE clinical guideline ***Bipolar disorder: assessment and management [CG185]***.

[Return to contents](#)

Parity of esteem

Parity of esteem is the principle by which mental health must be given equal priority and value to physical health. As a pharmacy professional, you are in an ideal position to work with those who have mental health conditions and their carers to ensure that patients receive equal treatment, whether it is for a mental or

physical condition. For more information about parity of esteem, access the NHS England blog on **Parity of esteem**.

[Return to contents](#)

Pharmacological treatment

All bipolar treatment should be initiated in secondary care. When symptoms have responded effectively to treatment, individuals should be offered the option to return to primary care for further management.¹⁰ The NICE clinical guideline ***Bipolar disorder: assessment and management [CG185]***, which we will refer to throughout this section, offers recommendations on initial management in secondary care.

Different pharmacological treatments can be used to help relieve symptoms associated with both of the elements of bipolar disorder: hypomania and mania, and depression.

Mania and hypomania

NICE clinical guideline ***Bipolar disorder: assessment and management [CG185]*** recommends treatment options for those who develop mania or hypomania in Section ***1.5 Managing mania or hypomania in adults in secondary care***.

Advance statements

Section 1.5 of NICE ***Bipolar disorder: assessment and management [CG185]*** advises to *‘Encourage people with bipolar disorder to develop advance statements while their condition is stable, in collaboration with their carers if possible.*

For information about advance statements and the difference between advance statements and advanced decisions, visit the Rethink Mental Illness ***Planning for your care – Advance statements and advance decisions*** page.

Depression

NICE’s ***Bipolar disorder: assessment and management [CG185]*** recommends treatment options for both those who are not already taking medicine to treat their bipolar disorder and those who are, in section ***1.6 Managing bipolar depression in adults in secondary care***.

A key section of this guideline for pharmacy professionals is ***1.10 How to use medication***. This section contains information on starting and monitoring antipsychotics in general and specific information relating to lithium, valproate and lamotrigine.

Lithium and sodium valproate are medicines where there are differences in equivalent doses of oral formulations. The Specialist Pharmacy Service (SPS) document ***Which medicines require extra care when switching between liquid and tablet/capsule formulations?*** considers which medicines may require dose adjustment when switching between oral formulations and presents information on dose adjustment in a summary table.¹¹

Visit the SPS information page, **Lithium monitoring**, for advice on the safe use of lithium, including monitoring and advice for patients.

Women and pregnancy

There is guidance on the use of valproate in women of childbearing potential and during pregnancy. Further information can be found on the Medicines and Healthcare products Regulatory Agency (MHRA) guidance for **Valproate use by women and girls** page.

College of Mental Health Pharmacists (CMHP) leaflets

The CMHP has produced four leaflets which look at antipsychotics, anxiety, antidepressants/depression and mood stabilisers which can be found in the CPPE mental health hub under **Mental health hub leaflets**. Each leaflet contains information about pharmacological treatment options and their indications, pharmacology, side effects (and their management), monitoring, interactions, referral information and key information to offer patients and their carers. Although these leaflets are aimed at community pharmacy, the information that they contain can be used in all areas of practice.

[Return to contents](#)

Non-pharmacological treatment

Non-pharmacological treatments for treating an episode of depression include psychological treatments similar to those offered to people with depression. NICE clinical guideline **Bipolar disorder: assessment and management [CG185]** recommends a psychological intervention that has been developed specifically for bipolar disorder or a high-intensity psychological intervention.¹²

High-intensity psychological interventions include cognitive behavioural therapy (CBT) – which looks at ways to change negative patterns in feelings, thoughts and behaviour, interpersonal therapy which focuses on relationships with others and how these relationships relate to feelings, thoughts and behaviour – and behavioural couples therapy (BCT), which looks at how to recognise and resolve emotional problems between partners.^{12,13} These high-intensity psychological interventions are also referred to as talking therapies.

During an episode of mania or hypomania, psychological treatments are not usually offered.¹³ NICE recommends that *‘people with mania or hypomania have access to calming environments and reduced stimulation. Advise them not to make important decisions until they have recovered from mania or hypomania and encourage them to maintain their relationships with their carers if possible.’*¹⁴

If people are experiencing a long and severe period of depression or a long period of mania and other treatments have not worked, or the situation is life-threatening, electroconvulsive therapy may be considered.¹⁵ Mind has a page, **Electroconvulsive therapy (ECT)**, which provides information on this potentially controversial treatment.

Visit Mind's ***Bipolar disorder – What treatments can help?*** page for an overview of talking therapies and longer-term management.

[Return to contents](#)

Patient support

Mind is a charity that provides advice and support to empower anyone experiencing a mental health problem. The following video by Mind – ***We are Mind*** – explores the work that Mind does to support people with mental health problems.



Bipolar UK is the national charity dedicated to supporting individuals with bipolar disorder and their families and carers. Bipolar UK offers support groups that can be found on the ***Find a peer support group*** page, a ***support line***, an ***eCommunity*** and ***work and learning***.

Resources for people with bipolar disorder

The Royal College of Psychiatrists' ***Learning disabilities*** page contains easy-to-read leaflets on a variety of topics including bipolar disorder, depression and medicines commonly used by those with bipolar disorder.

[Return to contents](#)

Further resources

Visit CPPE's ***Mental health*** gateway page for further learning on mental health in general.

NICE guideline ***Mental health problems in people with learning disabilities: prevention, assessment and management [NG54]*** 'covers preventing, assessing and managing mental health problems in people with learning disabilities in all settings (including health, social care, education, and forensic and criminal justice). It aims to improve assessment and support for mental health conditions, and help people with learning disabilities and their families and carers to be involved in their care.'¹⁶

Read the *British Medical Journal* clinical review article for further details about bipolar disorder and its management:

Anderson I, Haddad P, Scott J. ***Bipolar disorder***. *British Medical Journal*. 2012; 345:e8508.

Access the *Mental health awareness* e-learning via the **e-Learning for Healthcare learning modules** page on CPPE's website.

Read the Progress in neurology and psychiatry review article, Sie M, **Mood stabilisers in the management of bipolar affective disorder**. *Progress in neurology and psychiatry journal*. 2014;18.03.

The following studies explore the use of neuroimaging to try and determine the changes in the brain of someone with bipolar disease in order to try and determine its pathophysiology:

- Kempton MJ, Salvador Z, Munafò MR, et al. **Structural Neuroimaging Studies in Major Depressive Disorder: Meta-analysis and Comparison With Bipolar Disorder**. *Archives of General Psychiatry*. 2011;68(7):675-90.
- Adler CM, DelBello MP, Jarvis K, et al. Voxel-Based Study of Structural Changes in First-Episode Patients with Bipolar Disorder. *Biological Psychiatry*. 2007;61(6):776-81
- Hibar DP, Westlye LT, Doan NT, et al. **Cortical abnormalities in bipolar disorder: an MRI analysis of 6503 individuals from the ENIGMA Bipolar Disorder Working Group**. *Molecular Psychiatry*. 2018;23(4):932-42.
- Selvaraj S, Arnone D, Job D, et al. **Grey matter differences in bipolar disorder: a meta-analysis of voxel-based morphometry studies**. *Bipolar Disorders*. 2012;14(2):135-45.
- Phillips ML, Ladouceur CD, Drevets WC. **A neural model of voluntary and automatic emotion regulation: implications for understanding the pathophysiology and neurodevelopment of bipolar disorder**. *Molecular Psychiatry*. 2008;13(9):829, 833-57.
- Kraguljac NV, Reid M, White D, et al. **Neurometabolites in schizophrenia and bipolar disorder –A systematic review and meta-analysis**. *Psychiatry Research*. 2012;203(2-3):111-25.

[Return to contents](#)

External websites

CPPE is not responsible for the content of any non-CPPE websites mentioned on this page or for the accuracy of any information to be found there.

All web links were accessed on 3 February 2023.

[Return to contents](#)

References

1. The National Institute for Health and Care Excellence. Clinical Knowledge Summaries. **Bipolar disorder. Definition**. October 2022.
2. NHS. **Bipolar disorder. Overview**. January 2023.
3. NHS. **Bipolar disorder. Symptoms**. January 2023.

4. The National Institute for Health and Care Excellence. Clinical Knowledge Summaries. **Bipolar disorder. Causes.** October 2022.
5. Mind. **Bipolar disorder. Causes.** January 2023.
6. The National Institute for Health and Care Excellence. Clinical Knowledge Summaries. **Bipolar disorder. What is the prognosis?** October 2022.
7. BMJ. Best Practice. **Bipolar disorder in adults. Complications.** December 2020.
8. The National Institute for Health and Care Excellence. Clinical Knowledge Summaries. **Bipolar disorder. Complications.** October 2022.
9. The National Institute for Health and Care Excellence. Clinical guideline 185: **Bipolar disorder: assessment and management.** February 2020.
10. The National Institute for Health and Care Excellence. Clinical guideline 185: **Bipolar disorder: assessment and management. 1.9 Promoting recovery and return to primary care.** February 2020.
11. Specialist Pharmacy Service. **Which medicines require extra care when switching between liquid and tablet/capsule formulations?** 19 July 2021.
12. The National Institute for Health and Care Excellence. Clinical guideline 185: **Bipolar disorder: assessment and management. 1.6 Managing bipolar depression in adults in secondary care.** February 2020.
13. Mind. **Bipolar disorder. Treatments.** February 2022.
14. The National Institute for Health and Care Excellence. Clinical guideline 185: **Bipolar disorder: assessment and management. 1.5 Managing mania or hypomania in adults in secondary care.** February 2020.
15. The National Institute for Health and Care Excellence. Technology appraisal guidance 59: **Guidance on the use of electroconvulsive therapy.** 1 October 2009.
16. The National Institute for Health and Care Excellence. NICE guideline 54: **Mental health problems in people with learning disabilities: prevention, assessment and management.** September 2016.

[Return to contents](#)

Last review: March 2023

Next review due: March 2024